fl.

FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (REV. 11,2000)

TRANSMITTAL LETTER TO THE UNITED STATES

ATTORNEY'S DOCKET NUMBER 514413-3911

U.S. APPLICATION NO. (If known see 37 C.F.R. 1.5)

DESIGNATED/ELECTED OFFICE (DO/EO/US) N49410 CONCERNING A FILING UNDER 35 U.S.C. 371 INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE

PCT/EP00/07339 28 JULY 2000 9 AUGUST 1999

TITLE OF INVENTION HERBICIDAL COMPOSITIONS COMPRISING POST-EMERGENCE HERBICIDES FOR SOIL APPLICATION APPLICANT(S) FOR DO/EO/US Dr. Udo BICKERS and Dr. Gerhard FRISCH Applicants herewith submit to the United States Designated/Elected Office (DO/EO/US) the following items and other information: This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. This is an express request to promptly begin national examination procedures (35 U.S.C. 371(ft)). 4. The US has been elected by the expiration of 19 months from the priority date (PCT Article 31). 5. A copy of the International Application as filed (35 U.S.C. 371(c)(2)) is attached hereto (required only if not communicated by the International Bureau). has been communicated by the International Bureau. has been communicated by the international Busicalian is not required, as the application was filed in the United States Receiving Office (RO/US). An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). 7. Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))

are attached hereto (required only if not communicated by the International Bureau).

have been communicated by the International Bureau.

☐ have not been made; however, the time limit for making such amendments has NOT expired.
☐ have not been made and will not be made.

A English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).

 An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 20 below concern document(s) or information included:

11. An Information Disclosure Statement under 37 CFR 1.97 and 1.98.

12. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. A FIRST preliminary amendment.

 A SECOND or SUBSEOUENT preliminary amendment. 15. A substitute specification.

16. A change of power of attorney and/or address letter.

17. A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 18. A second copy of the published international application under 35

19. A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).

20. Other items or information:

U.S.C. 154(d)(4).

PCT/RO/101, PCT/IB/308, PCT/ISA/210 l page Abstract

EXPRESS MAIL

Mailing Label Number: EV001577608US Date of Deposit: February 7, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service

"Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents and Trademarks, Box PCT Washington, DC 20231.

(Signature of person mailing paper or fee)

The second second section is the second seco

U.S. APPLICATION	049410	0) INTERNAT P	IONAL APPLICATION NO. CT/EP00/07339			EY'S DOCKET NO. 14413-3911
21. A The follow	wing fees are submitted			(CALCULATIONS	PTO USE ONLY
Neither international pr nor international search	FEE (37 CFR 1.492) eliminary examination fee fee (37CFR 1.445(a)(2) p h Report not prepared by	(37 CFR 1.482) said to USPTO	\$1040.00			
	y examination fee (37 C.I al Search Report prepared		\$890.00			
International preliminar international search fee	y examination fee (37 CF (37 CFR 1.445(a)(2)) pai	R 1.482) not paid to USP d to USPTO	TO but\$740.00			
International preliminal but all claims did not sa	y examination fee paid to itisfy provisions of PCT A	USPTO (37 CFR 1.482) rticle 33(1)-(4	\$710.00			
International preliminar and all claims satisfied	y examination fee paid to provisions of PCT Article	USPTO (37 CFR 1.482) 33(1)-(4)	\$100.00			
ENTER APPROI	RIATE BASIC FE	E AMOUNT =		\$	890.00	
	of for furnishing the oat iest claimed priority da		nan 20 30	\$		
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$		
Total Claims	<u>13</u> - 20 =	<u>0</u>	x \$18.00	\$	00.00	
Independent Claims	<u>1</u> - 3 =	<u>0</u>	x \$84.00	\$	00.00	
MULTIPLE DEPEN	DENT CLAIM(S) (if a	pplicable)	+ \$280.00	\$	*	
		OF ABOVE CAL		\$	890.00	
Applicant claims above are reduced by	small entity status. Se	ee 37 C.F.R. 1.27. The	e fees indicated +	s		
			SUBTOTAL =	\$		
	30.00 for furnishing the		ter than 🔲 20 🔲 30	s		
months from the earl	iest claimed priority da		TIONAL TEEL -	S	890.00	······
Fee for recording the er	nclosed assignments (37 C		TIONAL FEE =	_		
	ropriate cover sheet (37 C	\$	40.00			
		TOTAL FEE:	S ENCLOSED =	\$	930.00	
				L	nount to be refunded:	S
				<u> </u>	charged:	\$
	in the amount of \$9					_
	ge my Deposit Accou copy of this sheet is		amount of \$t	o cc	over the above	tees.
c. The Commi overpaymen	ssioner is hereby aut at to Deposit Accoun	horized to charge an t No. <u>50-0320</u> . A du	y additional fees wh aplicate copy of this	ich she	may be require et is enclosed.	ed, or credit any
	oe charged to a credi nation should not be 38.					
NOTE: Where an : 1.137(a) or (b)) mus	appropriate time limi at be filed and granted	t under 37 CFR 1.494 I to restore the applic	or 1-495 has not been ation to pending state	1 m	et, a petition to	revive (37 CFR
SEND ALL CORRE	SPONDENCE TO:	_	Slu 1 (U	e	el_	
WILLIAM S. FRO			S.C. ATOKE			
	RENCE & HAUG LL		WILLIAM F. LAWR	EN	CE	
745 FIFTH AVENU NEW YORK, NEW			NAME			
			28,029			
Dated: February 7, 2	2002	_	REGISTRATION NU	MP	ER	

10/049410 JC11 Rec'd PCT/PTO 0 7 FEB 2002

514413-3911

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:

BICKERS, et al.

Filed .:

Filed Concurrently Herewith

Title of Invention:

HERBICIDAL COMPOSITIONS COMPRISING POST-EMERGENCE HERBICIDES FOR SOIL APPLICATION

> 745 Fifth Avenue New York, NY 10151

EXPRESS MAIL

Mailing Label Number: EV001577608US

Date of Deposit:

February 7, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Honorable Commissioner of Patents and Trademarks, Washington, DC 20231

(Signature of person mailing paper or fee)

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents

Box PCT

Washington, D.C. 20231

Sir:

Before the issuance of the first Office Action, please amend the above-identified

application as follows:

IN THE SPECIFICATION:

Please replace the paragraph beginning at page 2, line 26, with the following rewritten paragraph:

In the field of crop protection and, accordingly, also in the field of the control of harmful plants, there have been repeated attempts to enhance the use of properties of a given active compound, such as, for example, its activity spectrum, its persistence or the required application rate, for example by modified formulations or application methods. Thus, Weed Research, 1997, 37, 19-36, examines whether the activity of the herbicide chlorsulfuron can be enhanced and its tendency to leach can be reduced by using a controlled0release formulation. US 5,674,519 discloses that the tendency of certain crop protection agents including herbicides to leach in the soil can be reduced by a formulation in which the active compounds are present in microencapsulated form. Similarly, WO 98/05205 and US 5,543,383 describe a particular form of encapsulating crop protection agents which leads to an increased activity and a reduced susceptibility to leaching by rain. Furthermore, WO 99/26474 describes methods for releasing active compounds using cyclodextrins or carbohydrates. EP-A 0 517 669 discloses that the leaching behavior of agrochemicals can be improved by microencapsulation in polyester polymers. GB-A 1 041 028 provides mixtures of paraquat salts and fuller's earth having anticoccidial properties.

Please replace the paragraph beginning at page 3, line 8, with the following rewritten paragraph:

Accordingly, it is an object of the present invention to make possible the use of post-emergence herbicides by the pre-emergence method. This object is achieved by herbicidal compositions, comprising an effective amount of one or more post-emergence herbicides and an amount of a carrier material from the group consisting of fullers earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof, with

the proviso that herbicidal compositions comprising paraquat and fuller's earth shall be excluded.

Please replace the paragraph beginning at page 3, line 27, with the following rewritten paragraph:

Usually, the compositions according to the invention comprise

- a) from 0.15 to 48% by weight of one or more post-emergence herbicides,
- b) from 2 to 90% by weight of a carrier material and
- c) from 0 to 97% by weight of a solvent.

IN THE CLAIMS:

Please amend claim 1 as follows:

(Amended) A herbicidal composition, comprising an effective amount of one or more
post-emergence herbicides and an amount of a carrier material selected from the group
consisting of fuller's earth, aerogels, high-molecular-weight polyglycols and polymers based on
acrylic acid, methacrylic acid and copolymers thereof, with the proviso that herbicidal
compositions comprising paraquat and fuller's earth shall be excluded.

REMARKS

The Specification and Claim 1 has been amended to reflect the amendments made under PCT Article 34. Attached hereto is a marked up version of the changes made by the current amendment. The attached page is captioned "Version with markings to show changes made."

The filing fee has been calculated based upon the amendment made to the claim 1.

Respectfully submitted,

FROMMER LAWRENCE & HAUG LLP Attorneys for Applicant

By:

William F. Lawrence

Reg. No. 28,029

Tel. (212) 588-0800

JC11 Reg'd PCT/PTO 0 6 FEB 2002 514413-3911

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

Paragraph beginning at line 26 of page 2 has been amended as follows:

In the field of crop protection and, accordingly, also in the field of the control of harmful plants, there have been repeated attempts to enhance the use of properties of a given active compound, such as, for example, its activity spectrum, its persistence or the required application rate, for example by modified formulations or application methods. Thus, Weed Research, 1997, 37, 19-36, examines whether the activity of the herbicide chlorsulfuron can be enhanced and its tendency to leach can be reduced by using a controlled0release formulation. US 5,674,519 discloses that the tendency of certain crop protection agents including herbicides to leach in the soil can be reduced by a formulation in which the active compounds are present in microencapsulated form. Similarly, WO 98/05205[,too,] and US 5,543,383 describe a particular form of encapsulating crop protection agents which leads to an increased activity and a reduced susceptibility to leaching by rain. Furthermore, WO 99/26474 [and WO 99/26743] describes methods for releasing active compounds using cyclodextrins or carbohydrates. EP-A 0 517 669 discloses that the leaching behavior of agrochemicals can be improved by microencapsulation in polyester polymers. GB-A 1 041 028 provides mixtures of paraquat salts and fuller's earth having anticoccidial properties.

Paragraph beginning at line 8 of page 3 has been amended as follows:

Accordingly, it is an object of the present invention to make possible the use of post-emergence herbicides by the pre-emergence method. This object is achieved by herbicidal compositions, comprising an effective amount of one or more post-emergence herbicides and an amount of a

carrier material from the group consisting of fullers earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof, with the proviso that herbicidal compositions comprising paraquat and fuller's earth shall be excluded.

Paragraph beginning at line 27 of page 3 has been amended as follows:

Usually, the compositions according to the invention comprise

- a) from [0.001] <u>0.15</u> to 48% by weight of one or more post-emergence herbicides,
- b) from 2 to 90% by weight of a carrier material and
- c) from 0 to 97% by weight of a solvent.

In the Claims:

Claim 1 has been amended as follows:

(Amended) A herbicidal composition, comprising an effective amount of one or more
post-emergence herbicides and an amount of a carrier material selected from the group consisting
of fuller's earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid,
methacrylic acid and copolymers thereof, with the proviso that herbicidal compositions
comprising paraguat and fuller's earth shall be excluded.

Description

5

10

15

20

25

35

Herbicidal compositions comprising post-emergence herbicides for soil application

1

The invention relates to the technical field of herbicides, and in particular of herbicides for soil application.

For controlling undesirable harmful plants, the user can choose from a large number of herbicides which can be applied depending on the biological properties of the herbicides, the type of harmful plants to be controlled and the type of useful plants. Here, it has also to be taken into account that numerous herbicides, owing to their intrinsic mechanism of action, can be used either exclusively by the pre-emergence method or exclusively by the post-emergence method. Both methods and the herbicides used for these methods have certain advantages and disadvantages. Disadvantages of the post-emergence method where, for example, herbicides with foliar action such as bilanafos, diquat, glufosinate-ammonium, glyphosate and paraquat are also used, are considered to be, for example:

- damage to useful plants by overdosage
- depending on the type of herbicide used, on the harmful plants to be controlled and/or the useful plants in question, a need to repeat the application several times.
- The last mentioned disadvantage is particularly grave, especially from the point of view of time-economic farming.

On the other hand, numerous herbicides which can be used by the postemergence method have considerable advantages, in particular also from
an ecological point of view, since in many cases they have more favorable
toxicological and ecotoxicological properties than the herbicides which can
be used by the pre-emergence method. Depending on the particular
application situation, in many cases it would therefore be desirable for the
user to be able to use a post-emergence herbicide under the conditions of
a pre-emergence application, i.e. prior to the emergence of the plants or, if
appropriate, simultaneously with the sowing of the useful plants. However,
hitherto such an application has, in particular in the case of herbicides with
foliar action, not been possible, for reasons of the mechanism of action on

which these herbicides are based, since herbicides with foliar action in the form of the herbicidal compositions of the prior art, for example, are only taken up via the green parts of the plant, for reasons of their leaching properties or else for reasons of their degradation behavior in the soil. Glufosinate-ammonium (2-amino-4-(hydroxymethylphosphinyl)butanoic acid), for example, which can be used as herbicide with foliar action is known to be decomposed rapidly in the soil, so that it is not capable of displaying any herbicidal action (G. Hoerlein in "Reviews of Environmental Contamination and Toxicology", vol. 138, Springer-Verlag; "The Pesticide Manual", 11th Edition, 1997, British Crop Protection Council). The herbicide glyphosate (N-(phosphonomethyl)glycine), which likewise has foliar action, is known to be adsorbed strongly by the soil and to be degraded therein, so that it is not available to the plant in sufficient amounts (L. Torstensson in "The Herbicide Glyphosate", Butterworths, pp. 137-150).

Hereinbelow, the term "herbicide" is, depending on the context, to be understood as meaning the pure herbicidally active compound or the herbicidally active compound in formulated form, i.e. the herbicidal composition. The terms "pre-emergence" and "post-emergence" are here to be understood as referring to the time of the emergence of the harmful plants, i.e. a treatment by the pre-emergence method means a treatment with a herbicide prior to the emergence of the harmful plants and thus prior to the formation of green parts of plants. Analogously, a treatment by the post-emergence method means a treatment with a herbicide after emergence of the harmful plants.

In the field of crop protection and, accordingly, also in the field of the control of harmful plants, there have been repeated attempts to enhance the use properties of a given active compound, such as, for example, its activity spectrum, its persistence or the required application rate, for example by modified formulations or application methods. Thus, Weed Research, 1997, 37, 19-26 examines whether the activity of the herbicide chlorsulfuron can be enhanced and its tendency to leach can be reduced by using a controlled-release formulation. US 5,674,519 discloses that the tendency of certain crop protection agents including herbicides to leach in the soil can be reduced by a formulation in which the active compounds are present in microencapsulated form. WO 98/05205, too, describes a particular form of encapsulating crop protection agents which leads to an increased activity and a reduced susceptibility to leaching by rain.

10

15

20

25

35

Furthermore, WO 99/26474 and WO 99/26743 describe methods for releasing active compounds using cyclodextrins or carbohydrates.

However, none of the publications mentioned indicates that it may be possible to use post-emergence herbicides pre-emergence by employing suitable measures.

Accordingly, it is an object of the present invention to make possible the use of post-emergence herbicides by the pre-emergence method. This object is achieved by herbicidal compositions, comprising an effective amount of one or more post-emergence herbicides and an amount of a carrier material from the group consisting of fullers earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof.

Here, the compositions according to the invention should only comprise post-emergence herbicides which, in the form of the herbicidal compositions of the prior art, can be used exclusively by the post-emergence method, i.e. after emergence of the undesirable harmful plants.

Surprisingly, post-emergence herbicides which are present bound to the carrier materials mentioned above display herbicidal action against undesirable harmful plants when used by the pre-emergence method, i.e. when the herbicidal compositions have been applied prior to the emergence of the harmful plants.

Usually, the compositions according to the invention comprise

- a) from 0.001 to 48% by weight of one or more post-emergence herbicides.
- 30 b) from 2 to 90% by weight of a carrier material and
 - c) from 0 to 97% by weight of a solvent.

Particularly suitable post-emergence herbicides are those from the group of the herbicides with foliar action. Preference is given to herbicides from the group consisting of bilanafos, diquat, glufosinate-ammonium, glyphosate and paraquat. Particular preference is given to glufosinate-ammonium. The abovementioned herbicides are known, for example, from "The Pesticide Manual", 11th Edition, 1997, British Crop Protection Council. The herbicides in question can, of course, also be employed in the commercial

salt form, as known, for example, from "The Pesticide Manual".

It is a further advantage of the compositions according to the invention that they are present in solid form and can be used, for example, in the form of granules. The user can apply this solid presentation form directly to the area to be treated, i.e. without it being necessary to prepare a spray liquor. To this end, they are, depending on the type of useful plants and the type of expected harmful plants to be controlled, applied onto the soil to be treated, worked into the soil or applied by side application.

Side application means that the herbicide is applied in the soil below the seeds. Here, it is particularly advantageous that the application of the herbicidal compositions can also take place in the same operation as the sowing of the crop plants, resulting in the soil being subjected to reduced mechanical stress.

15

20

25

10

5

The herbicides which can be used for the compositions according to the invention can be processed to the compositions according to the invention in the form of the pure active compound or else in the customary commercial formulation together with carrier materials and, if appropriate, additional substances, in the manner described further below. A particular advantage of the compositions according to the invention is their good environmental compatibility. Owing to the fact that the herbicides, in contrast to conventional application methods, are not sprayed in liquid form but applied in solid form, there is no risk of uncontrolled drift of spray mists onto the user and adjacent areas and plants. Moreover, environmentally compatible formulation auxiliaries are used as they are either - like the high-molecular-weight polyglycols and polymers of acrylic acid, methacrylic acid and copolymers thereof - biodegradable to unobjectionable substances or - like fuller's earth - show environmentally neutral behavior.

30

35

The herbicidal compositions are prepared by mixing the active compounds, carrier materials and, if appropriate, further additives in question according to methods known to the person skilled in the art. Granules, for example, can be prepared by the methods described in EP-A - 0 413 267. Depending on the carrier material, the substances in question can be present in solid or liquid or dissolved form. In the preparation of granules, it is advantageous for the active compounds and additives to be incorporated to be present in liquid or dissolved form. Here, these substances can be applied uniformly in the known manner for example by spraying, pouring,

dropwise addition, treatment in a fluidized bed, concrete mixer, tumble mixer, etc. It is also possible to oversaturate these carrier materials with solutions of the active compounds and additives in question to form, for example, gels, or to remove excess water by drying. In the case of the preparation of melt granules, for example melt granules based on polyethylene glycol, active compounds and additives are preferably incorporated in solid form and extruded. Also possible is tabletting, pelletizing, the preparation of flakes and comminution to powders by breaking or grinding. In all preparation processes, aftertreatment by comminution may take place. It is also possible to compact very fine granules using suitable adhesives to avoid the formation of dust. It is furthermore possible, if appropriate, to add additives to improve, for example, the flowability of the granules or their wettability.

Depending on the intended purpose, the herbicidal compositions may additionally comprise other substances which can be used in crop protection, such as pre-emergence herbicides, plant growth regulators, fungicides, insecticides, safeners ((herbicidal) antidotes), nutrients, preservatives, seed dressings and fertilizers. The selection of the pre-emergence herbicides, plant growth regulators, fungicides, insecticides and seed dresssings to be added, if appropriate, depends on the type of useful plants and the type of expected harmful plants, fungi, insects and feed pests to be controlled. In principle, it is possible to use all commercial substances which are usually employed in crop protection. Suitable nutrients and fertilizers are, in particular, aqueous ammonium nitrate/urea solutions and NPK solutions such as 12-6-8, 8-8-6, 5-8-10, and also ammonium sulfate and/or ammonium nitrate solutions.

Preferred carrier materials are aerogels, polymers based on acrylic acid, methacrylic acid and copolymers thereof and also high-molecular-weight polyglycols.

When using the compositions according to the invention, it has to be ensured that the useful plant is sufficiently tolerant to the herbicide 35 employed. In addition to useful plants in which sufficient tolerance to herbicides has been achieved by targeted genetic manipulation, this includes plants which, such as, for example, barley, have sufficient tolerance to some herbicides even without genetic manipulations. The compositions according to the invention can be used particularly

advantageously in crops of useful plants from the group consisting of cereals, corn, soyabean and rapeseed. They are preferably used in crops which are sufficiently tolerant to the herbicide in question owing to genetic manipulation.

5

10

15

20

25

30

35

As in the case of other herbicidal compositions, it is, of course, in the case of the compositions according to the invention also possible and in some cases advantageous to provide the herbicide with additives such as surfactants, wetting agents, emulsifiers, adjuvants, ammonium salts, preservatives, colorants, antifoams, tackifiers, solvents, buffer systems and UV stabilizers. Depending on the type and purpose of the additives, these can be processed together with the herbicide in a formulation or else be present and subsequently used separately from the herbicide. These additives serve to improve application properties. The use of such additives may be advantageous to enhance, for example, the persistence of the herbicide. Preservatives are used, for example, to slow down the biological degradation of active compounds and/or formulation auxiliaries.

Suitable surfactants are, for example, condensates of formaldehyde and phenol and/or naphthol, in each case with or without sodium bisulfite, such as Tamol NNO from BASF. Rapidamin-Reserve C from Clariant or Galoryl MT800 or DT201 from CFPI: C12-C24 fatty alcohols having 2 to 40 EO and/or PO, if appropriate phosphated and/or neutralized with alkanolamine, alkali metal or ammonia; di- and tristyrylphenyl analogs of the fatty alcohols mentioned above, such as, for example, the Genapol series from Clariant, Grafol types from Henkel and Soprophor types from Rhodia; alkyl ether sulfates such as Genapol LRO® from Clariant; alkyl- and/or alkenylsulfonates such as Hostapur OS from Clariant, ligninsulfonates such as Ufoxane 3A and Vanisperse CB from Booregard; Reax types from Westvaco; N-methyltaurides such as Hostapon T from Clariant: sulfosuccinic monoester salt such as Hoe S 1728 from Clariant; alkyl polysaccharides such as Plantaren APG 600 from Henkel; ethoxylated C12-C24-fatty amines, such as the Genamin types from Clariant. Suitable solvents are, in each case depending on the intended purpose, alcohols, diols, polyols, N-substituted pyrrolidones, ketones, aldehydes, ethers, polyethers, paraffins, aromatic compounds, heteroaromatic compounds, cycloalkanones, dimethylsulfoxide, tetrahydrofuran and water. It is advantageous to employ water.

25

30

5

Suitable preservatives are, for example, Bronidox L from Henkel, Mergal types from Riedel de Haen, Proxel from ICI, ascorbic acid derivatives, benzoic acid derivatives, formaldehyde, citric acid, preservatives from the Kathon series from Rohm & Haas and brohopol from BASF.

Suitable adjuvants are, for example, alkyl polysaccharides or lauryl ether sulfates.

Suitable carrier materials from the group of the high-molecular-weight polyglycols are, in particular, polyethylene glycols having a molecular weight of from 2000 to 40,000 (PEG 2000 to PEG 40,000). Suitable carrier materials from the group of the aerogels are, in particular, the aerogels described in EP-A 0 171 722.

15 It is particularly advantageous to add a substance such as ammonium sulfate, ammonium nitrate and one of the surfactants mentioned above, in particular a surfactant from the group of the alkyl ether sulfates.

The use of the herbicidal compositions by the pre-emergence method is novel and also forms part of the subject matter of the invention.

Using the herbicidal compositions according to the invention, it is possible to control undesirable harmful plants by the pre-emergence method. This method is new and also forms part of the subject matter of the invention.

The invention is illustrated by the embodiments that follow.

A. Formulation examples

By way of example, the qualitative and quantitative (in % by weight) composition of numerous herbicidal compositions according to the invention is listed in Tables 1 to 4.

l able 1	Herbicide (active compound)	(punoduoo	Carrier materials				Additives
Ö.	sasta [®] 50% (glu- ferialso)	etsed® 1/g 02t -ulg) (etsnisot	Luquasorb AF1	droasorb AF2	Aerogel 889	Aerogel	water
-	2			06			8 00
2	2			10			88.00
8	2			4			94.00
4	2			2			96.00
2	2		06				8.00
9 1	2		88				10.00
,	2		20	*			48.00
80	2		10				88.00
6	2		4				94.00
10	2		2				96 00
-1		95	2				9
12		06	10				
13	06		10				
14	95		2				
15	10		06				
16	2				06		8.00
17	2				80		18.00
18	2				40		58 00
19	2				20		48.00
20	2				65		33.00
21	2					90	38.00
22	2					20	48.00
						3	2

Basta 150 g/l: 13.5% of glufosinate-ammonium, 58.81% of Genapol LRO, 10.0% of Dowanol PM, 0.25% of Fluowet PL80, 0.005% of Duasyn acid blue AE, water ad 100%

			8	63.00	8						Γ	T	T	T	T	1	T	٦	5.00	0	8	2	2	3 5	3	-
	vater	٨	83	8	88	4	1			L	L	L	L	ļ	1	1	1		15.	9.00	17.	14.00	14.5	3 5	-	
	hycerol	3												Ş	2					9						
	lio bəəsəqs	,															2								Ī	
	bropylene Stycol	3					-	5.89	5.99	5.99	1.99	42.95	39.25												6	
	Stepanol ME				1	1			2	-				Ī	Ī		1	1						İ		
	Silcolapse 5020											4	4		Ī	1	1	1							t	
Additives	Duasyn acid blue AE						,	0.0	0.0	0.01	0.01	0.05	0.05			1	1	1	1						T	
Adc	Sipernat 22S							9	ç	2	2						u	,			2	S	2	2	t	
	Genapol LRO												10		9	2										
	Geropon SDS						I					9	10													
	Hostapur OSB					Ī	Ľ	Ì												1						
	urea			Ī	Ī	Ī			1	-	ç	-	-													
	ammonium sulfate				Ī												Ī		I				-	4		
	ammonium nitrate						-	-	-									Ī	Ī	-	-	4				
S	Aerogel C373	40	35	90														Ī		Ī					_	-
Carrier materials	889 legoreA								T		4	9	9												_	
Carrier	Luquasorb AF1						84	2	98	3 8	8		,	n	2	22	84	2	ıc.	7,	1	2	9	72	_	
de	%88 sise8®				15	2					T													1	_	
Herbicide	@Basta 50%	2	2	2	82	92	2	2	0	10	10	4	7 20	S	82	82	Ξ	8	8	2	0	4	,	7		
T I	ou.	23	54	25	56	27	28	83	30	8	S	3 6	3 5	\$	33	36	37	38	39	40	41	,	2 9	3	44	

တ

				_	_	_	_		_			-	_	_	_	_								
		water	1.	16.70	12.80		16.99	15.99		000	9.6	0	9.99					20 00	30.00	30.00	38.00	38.00	38.00	23.00
		Hostapur 820						Ī											ç	2	1	1	Ī	
		T nogstapon T	I	I			L	I		I								ç	2	I			I	
		дгусегоі	- 1									100	*						ļ					
		Supragil WP								2	2							I	T	T	T		T	T
		alycol propylene							33 99	2000	2 00	6.00	0000	52.33	80.0	0.00	5.89		I					
		Silcolapse 5020	Ī	Ī			Ī		4	T		7	,	‡	1	T		T	T	T	1	1	T	T
		duasynacid Alue AE					0.01	0.01	0.01		0	000	5	500	5 6	5 6	0.0	I		Ī	T		T	
	Se	Mergal K9N									0.4	;			Ī	T			Ţ	T	Ī			
	Additives	Cobate C	Γ										I	I		I	o r	_	L	I	I	I	I	
	A	Kathon MK								L						5	5	L			1			L
		D xobinor L	L								ŀ	L			5	5								
		Proxel GIXL												5	3							T		Γ
		sipernate 22S	2	2	2	2	5	2			5			Ľ	2	2 1	2			Ī	Ī	T	Ī	T
		KE 640								L	I			Ī						Ι	I	I	ç	2
		Genamin 1150																		I	I	40	2	
		Genapol							10			9			Γ	T					ç	2	T	Γ
		Geropon							9			9	10	2	Ī	l		Γ		ç	1		T	Γ
		(NH4) ⁵ 2O ⁵						2					10	2					Γ		T			
		NHH4NO3	Ц						_	L	-		-	Ŀ	-	-	-				I		I	
	÷	Aerogel	Ш						4	45	L	8	40	L	L	L		22	22	50	3 6	3 6	2	45
	Carrier mat	Luquasorb				75					L													
	Carri		75	75	13		72	33		Г	88			88	88	88	98							
	(active	Grandup 360 (glyph- 1/p (alyphasid) 1/p (alyphasid)		3.3																				
	F	paraquat 200 o/l	5.5																		I	İ		
က	Herbicide compound)	Basta %021		į,	7:7									Ĺ	Ĺ	L								
Table 3	를 함	%02 stes€	Ц	1	-	-+	2	-	-		-	-	_	₽-	⊢	-		Н	ш	_	-	2	├-	7
-		.oN	9	4	8	\$	잆	5	12	23	5	55	28	57	28	23	9	61	82	83	18	18	99	67

				_		_	_,	_		_								_				
		vater		30.00	38.00	38.00	38.00	43.00	43.00	43.00	43.00	43.00	25.00	52.00	43.00	43.00	43.00	43.00	43.00	43.00		Ī
	кэи	Mergal	1			1	1			1					1		1				0.1	-
	S SS I	Sipema													1		1				2	u
	dW II	Supragi	- 1	T	Ī		1													ဝ		
	tsi	SGF Rhodon		Ī	I		l										1		9			Γ
	uəß	Emulso	Ī	l						T	Ī			T			,	2				
	hor F1	Soprop		I		Ī									T	,	2	1				
	Por 3	D33 Sobrop		Ī	T				T	1	1			T	,	2	T					
	26	Berol 9			I			Ī	T		I			,	2		Ī					
	HS0	Tylose	Ī		Ī	Ī					Ī		١,	-	T		T					
Additives	18-88	loiwoM	L	I		Ī	T						-		T	T				-		_
Adc	0 10	Genapi 080	I			Ī		T	Ī	Ī	,	2		Ī	T			Ī				-
	1	KBE Byosby			Ī	Ī		T	T	ç	2			Ī	T	T		Ī		1	1	_
		qstaoH 062A2		l	Ī				۶	2			T	Ī	T		T		1		1	
	lo	Genap						ç	2			T					T	Ī				
	982	ECD 4					ç	2									-					
	on K12	Техарс				2	1		L									-				
	dW lig	Suprag			10	L	L	L														
	TM i	Galory 800		9			L	L			l					L	L					
	IS inse		9		L		L			L												_
	uo	Geropi	L	L	L	L	L	L	L	L				L							ŀ	2
	T noc	Hostap	L	L		L	L				L					L	L				n	
ier	hylene 20,000																			6	88.0	82.9
Carrier mat.	889 le	Aerog	20	20	20	20	45	45	45	45	45	15	45	45	45	45	45	45	45	?	I	
Herbicide	%66 E	etes 6					L		L					L						-	-[-	n
Hert	%09 E	stas 8®	7	2	7	2	8	8	2	2	2	1	0	7	2	2	7	2	,	1	Ī]
1		ou.	88	68	2	7	2	73	4	13	9	1	180	62	8	٦	22	33	5	1	3 8	g

Toblo 4

0.05 00 0.0 0.1 Mergal K9N 0.0 quì Stepanol ME 0.05 0.00 0. 0.05 **BA** euld quasyn acid Additives 0.05 0.5 Bronidox L 5 8 8 nitrate ammonium 0.0 0.0 980 Hostapur 46.55 46.75 47.39 47.39 47.89 47.69 46.69 46.59 46.39 46.85 47.85 47.74 diycol bropylene 98.85 98.65 98.50 97.95 Carrier material **BIACOI 20,000** bojkethylene 0.04 6 Aerogel P88 2 0.5 2,4-d-DMA techn. grade AMG-q-qorq 0.5 0.2 qichloro-Grade o, 5 6 diuron techn. techn. grade κi 300 Herbicide enizemis Grade 0.3 CMPP techn. Table 1a - Combinations 1.05 1.05 1.05 1.05 1.05 1.05 1.05 recuur duage mono-linuron 1.0 0.3 848ta 99% 20 2002 2.0 2.0 2002 202 Basta 50% 110 988888888888 ou.

Fable 2a - Combinations

5 5 - - -27 5 0.2 5 Mergal K9N 0.0000 0.01 0.01 10.0 10.0 Stepanol ME dry 10.0 0.01 Geropon SDS 0.00 0. 0.05 Additives qnasyn acid blue 0.5 nitrate 90 5 5 5 5 000 0. 0.0 0. ammonium 0.0 Hostapur OSB 46.48 47.63 47.85 47.85 6.49 46.80 46.35 47.45 45.95 44.45 46.89 46.79 18.65 47.85 bropylene glycol 86.0 Luquasorb AF1 Carrier material 5.0 5.0 Sipernat 22 S glycol 20,000 98.0 93.9 93.2 98.0 polyethylene 0.000000 0.04 0.04 0.04 0.0 40.0 Aerogel P88 grade 9 atrazine techn. techn. grade Ξ amidosulfurone grade 0.5 pentazone techn. 0.05 cycocei grade recuu. linuron Herbicide techn. grade 0 acetochlorine techn. grade metamitron techn. grade 0.06 MCPA-DMA salt 2,4-d-DMA salt 0 0 Basta 99% 30.0 20 200 000 200 Basta 50% 14 115 9 ន 2 2 ou.

							_			
		gibberelic acid Na							5	0.0
		3-indolyl- acetic acid Na					-		0.50	
		1-naphthyl- -acetic acid Na			1		2			
		phenyl- acid	5.0	000	ì					
		K8N Weldsj	0.1	-	-	-			0.1	
	Additives	Stepanol ME Dry	10.0	10.0	10.0	10.0			10.0	200
	Add	Geropon SDS								
		duasyn acid blue AE	0.5	0.05	0.05	0.05	0.05	0.05	0.05	30.0
		Bronidox L					0.5	0.5		
		muinomms etstin	1.0	1.0	1.0	1.0	0.	1.0	1.0	-
		Hostapur OSB					10.0	10.0		
		alycol propylene	46.35	44.85	46.75	45.85	46.44	46.25	46.84	46.65
	Carrier material	Aerogel 889	40.0	40.0	40.0	40.0	40.0	40.0	40.0	40.0
	Herbicide	%02 stead	2.0	2.0	2.0	2.0	2.0	2.0	2.0	5.0
Table 3a		ou.	134	135	136	137	138	139	140	141

20

25

30

35

B Biological examples

Meanings of the abbreviations used below:

5 ABUTH Abutilon theophrasti AMARE Amaranthus retroflexus BRSNW Brassica napus CHEAL Chenopodium album GALAP Galium aparine HORVS Hordeum vulgare LOLMU Lolium multiflorum

10 B.1 Action against harmful plants of glufosinate-ammonium bound to a polymer carrier material, compared to that of glufosinateammonium applied by a conventional method

In a greenhouse, a granulous formulation, prepared according to example 48, of the herbicide glufosinate-ammonium was broadcast at an application rate of 500 g of active compound per hectare (converted) on humus-containing loam soil, and the soil was then watered. Three days after the application, seeds of Setaria viridis were sown at a depth of 1 cm. For the entire duration of the experiment, a daytime temperature of 22-24°C and a nighttime temperature of 16-18°C were maintained, a uniform day length of 16 hours being achieved by additional illumination with sodium vapor lamps (approximately 7000lux). The relative atmospheric humidity was 60-80%. Visual scoring 35 days after sowing showed that an activity of 70% against Setaria viridis was achieved. For comparison, glufosinate-ammonium was applied as an aqueous solution, under otherwise identical conditions and likewise at an application rate of 500 g of active compound per hectare (converted). Visual scoring 35 days after sowing showed that the activity achieved against Setaria viridis was nil (0%).

B.2 Comparison of the action against harmful and useful plants of glufosinate-ammonium bound to polymer carrier material

In a greenhouse, seeds of the harmful plants LOLMU, AMARE and CHEAL and the useful plant HORVS were sown on humus-containing loam soil, covered with a 0.5 cm layer of the same soil and then watered. One day after sowing, a formulation, prepared according to example 48, of the herbicide glufosinate-ammonium was broadcast on the soil at an application rate of 1000 g of active compound per hectare (converted). For the entire duration of the experiment, a daytime temperature of 22-24°C and a nighttime temperature of 16-18°C were maintained, a uniform day length of 16 hours being achieved by additional illumination with sodium

vapor lamps (approximately 7000 lux). The relative atmospheric humidity was 60-80%. Visual scoring 35 days after sowing gave the results shown in Table 5, and it was found that, in contrast to the three harmful plants tested, the crop plant HORVS had not been damaged by the herbicide.

5

10

15

25

Table 5		Activ	ity [%] agai	inst harmful	and useful	l plants
Treatment according example No.	with to	herbicide formulation	LOLMU	AMARE	CHEAL	HORVS
	48		40	70	60	0

B.3 Action against harmful plants of glufosinate-ammonium bound to different carrier materials

Seeds of the harmful plants GALAP, AMARE, CHEAL and LOLMU were sown outdoors on humus-containing loam soil, covered with a 0.5 cm layer of the same soil and then watered. One day after sowing, formulations, prepared according to examples No. 40, 45 and 46, of the herbicide glufosinate-ammonium were broadcast at an application rate of 750 g of active compound per hectare (converted) on the soil. Visual scoring 35 days after sowing gave the results shown in Table 6.

Table 6			Ac	ctivity [%] a	gainst harm	ful plants
Treatment	with	herbicide	GALAP	AMARE	CHEAL	LOLMU
according	to	formulation				
example No						
	40		80	10	15	25
	45		100	35	70	30
	56		70	40	85	25

20 B.4 Effect of different fertilizers on the efficacy of glufosinateammonium bound to polymer carrier material

In a greenhouse, seeds of the harmful plants ABUTH, AMARE, CHEAL and LOLMU were sown in humus-containing loam soil, covered with a 0.5 cm layer of the same soil and then watered. One day after sowing, a formulation, prepared according to examples 41 and 43, of the herbicide glufosinate-ammonium was broadcast at an application rate of 750 g of active compound per hectare (converted) on the soil. For the entire duration of the experiment, a daytime temperature of 22-24°C and a nighttime temperature of 16-18°C were maintained, a uniform day length of

15

20

16 hours being achieved by additional illumination with sodium vapor lamps (about 7000 lux). The relative atmospheric humidity was 60-80%. Visual scoring 35 days after sowing gave the results shown in Table 7.

5 Table 7 Activity [%] against harmful plants Treatment with herbicide ABUTH AMARE CHEAL LOLMU according to formulation example No. 41 80 90 85 15 43 75 75 75 10

B.5 Effect of different surfactants on the efficacy of glufosinateammonium bound to carrier materials

In a greenhouse, seeds of the harmful plants GALAP, AMARE, CHEAL and LOLMU were sown in humus-containing loam soil, covered with a 0.5 cm layer of the same soil and then watered. One day after sowing, a formulation, prepared according to examples 40, 29 and 28, of the herbicide glufosinate-ammonium was broadcast at an application rate of 750 g of active compound per hectare (converted) on the soil. For the entire duration of the experiment, a daytime temperature of 22-24°C and a nighttime temperature of 16-18°C were maintained, a uniform day length of 16 hours being achieved by additional illumination with sodium vapor lamps (about 7000 lux). The relative atmospheric humidity was 60-80%. Visual scoring 35 days after sowing gave the results which showed that the efficacy of the herbicide is enhanced by adding surfactants to glufosinate-ammonium bound to carrier materials, see Table 8.

Table 8 Activity [%] against harmful plants

Treatment with herbicide according to formulation example No.	GALAP	AMARE	CHEAL	LOLMU
40 (without surfactant)	80	10	15	25
28 (with surfactant)	98	60	98	60
29 (with surfactant)	90	60	90	40

25 B.6 Effect of preservatives on the efficacy of glufosinateammonium bound to polymer carrier material

In a greenhouse, seeds of the harmful plants ABUTH and BRSNW were sown in humus-containing loam soil, covered with a 0.5 cm layer of the

same soil and then watered. One day after sowing, a formulation, prepared according to examples 40, 54 and 60, of the herbicide glufosinate-ammonium was broadcast at an application rate of 750 g of active compound per hectare (converted) on the soil. For the entire duration of the experiment, a daytime temperature of 22-24°C and a nighttime temperature of 16-18°C were maintained, a uniform day length of 16 hours being achieved by additional illumination with sodium vapor lamps (about 7000 lux). The relative atmospheric humidity was 60-80%. Visual scoring 35 days after sowing gave the results which showed that the efficacy of the herbicide is enhanced by adding preservatives to glufosinate-ammonium bound to carrier materials. see Table 9.

Table 9	Activity [%] against harmful plants
Treatment with herbicide according for formulation example No.	ABUTH	BRSNW
40 (without preservative)	30	0
54 (with preservative)	55	10
60 (with preservative)	55	70

10

- A herbicidal composition, comprising an effective amount of one or more post-emergence herbicides and an amount of a carrier material selected from the group consisting of fullers earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof.
- 2. The herbicidal composition as claimed in claim 1, comprising
 - a) from 0.15 to 48% by weight of one or more herbicides,
 - b) from 2 to 90% by weight of a carrier material and
 - c) from 0 to 97% by weight of a solvent.
- 3. The herbicidal composition as claimed in claim 1 or 2, wherein the herbicide is from the group of the herbicides with foliar action.
 - The herbicidal composition as claimed in any of claims 1 to 3, wherein the herbicide is from the group consisting of bilanafos, diquat, glufosinate-ammonium, glyphosate and paraquat.
 - The herbicidal composition as claimed in any of claims 1 to 4,
 wherein the herbicide is glufosinate-ammonium.
- 6. The process as claimed in any of claims 1 to 5, wherein the carrier materials are from the group consisting of aerogels, high-molecularweight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof.
- The herbicidal composition as claimed in any of claims 1 to 6, which
 additionally comprises at least one substance from the group
 consisting of pre-emergence herbicides, plant growth regulators,
 fungicides, insecticides, safeners, nutrients, seed dressings and
 fertilizers.
- 35 8. The herbicidal composition as claimed in any of claims 1 to 7, which additionally comprises at least one additive from the group consisting of surfactants, wetting agents, emulsifiers, adjuvants, ammonium salts, preservatives, colorants, antifoams, tackifiers, solvents, buffer systems and UV stabilizers.

 The herbicidal composition as claimed in claim 8, wherein the additive is from the group consisting of Genapol LRO[®], ammonium sulfate and ammonium nitrate

- 10. The use of herbicidal compositions as claimed in any of claims 1 to 9 for controlling undesirable harmful plants, which comprises applying the compositions by the pre-emergence method.
- 10 11. The use of herbicidal compositions as claimed in claim 10, wherein the compositions are used in crops of useful plants which are tolerant to the herbicidally active compounds contained in the compositions.
- 15 12. The use of herbicidal compositions as claimed in claim 11, wherein the compositions are used in crops of genetically modified useful plants.
- 13. A method for controlling undesirable harmful plants, wherein the 20 post-emergence herbicides are used in the form of a herbicidal composition as claimed in any of claims 1 to 9 by the pre-emergence method.

1999/M 223 WO

Abstract:

Herbicidal compositions comprising post-emergence herbicides, carrier materials from the group consisting of fullers earth, aerogels, high-molecular-weight polyglycols and polymers based on acrylic acid, methacrylic acid and copolymers thereof and, if appropriate, further additives, which herbicidal compositions can be used pre-emergence for controlling undesirable harmful plants, are described.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

	INVENTOR(S) / Residence			
, 0	1) Dr. Udo Bickers, Südstraße 2, D-498			
	Signature: Uan Droce		Date:	12.12.10
3 . (2) Dr. Gerhard Frisch, Westerwaldstrat	ße 7, D-612 <u>73 Wehrhe</u> i		00 .
	Signature:full_tfs	>	Date:	12.12.01
	The inventors are citizens of Germany			
67 83 80	Post Office Address of all Inventors:	Aventis CropScieno	e GmbH	

Patent- und Lizenzabteilung Industriepark Höchst Gebäude K 801 D-65926 Frankfurt Germany

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Herbicidal compositions comprising post-emergence herbicides for soil application

the specification of which

- is attached hereto
- was filed on July 28, 2000 as International Application PCT/EP00/07339 and including all the amendments through the date hereof.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s) for which Priority is Claimed:

Federal Republic of Germany, 19936784.1 of August 9, 1999

And I hereby appoint

William F. Lawrence, Registration No. 28,029 of the firm FROMMER LAWRENCE & HAUG, LLP whose post office address is 745 Fifth Avenue, New York, New York 10151, or their duly appointed associate, my attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and specify that all communications about the application are to be directed to the following correspondence address:

William F. Lawrence, Esq. c/o FROMMER, LAWRENCE & HAUG LLP 745 Fifth Avenue New York, New York 10151 Direct all telephone calls to: (212) 588-0800, to the attention of: William F. Lawrence